Journal of **Clinical Nursing**

ORIGINAL ARTICLE

A survey of parental self-efficacy experiences: maximising potential through health visiting and universal parenting support

Karen A Whittaker and Sarah Cowley

Aims. To examine parental self-efficacy experiences for users of a parenting support programme and consider the pertinence of self-efficacy theory to health visiting (public health nursing) practice.

Background. Commonly, successful parenting training programmes are underpinned by social learning principles and aim to strengthen parental self-efficacy. However, research examining programme effectiveness rarely discusses how self-efficacy outcomes are achieved.

Design. A descriptive survey was completed as the first part of a realistic evaluation study examining how a UK parenting support programme worked.

Method. The first part of the realistic evaluation involved validating outcome measures (the Parenting Self-Agency Measure and Self-Efficacy for Parenting Tasks Index subscales) and administering a questionnaire survey. The questionnaire was completed by adults accessing a parenting support programme during a 10-month period (n = 168). Data were analysed using descriptive and inferential statistics.

Results. Women were the main users of the programme, which included informal drop-in groups as well as more formalised health visiting services and parenting training courses. The Parenting Self-Agency Measure results indicated good general parental self-efficacy; however, the task-specific Self-Efficacy for Parenting Tasks Indexes scales suggested that parents were less self-efficacious in disciplining children. Lower self-efficacy scores correlated with high ratings for 'feeling tired', 'receiving negative comments' and 'giving-in to a child's demands'.

Conclusions. Study results indicate that the domain general and task-specific measures provide different, but helpful, insights into parental self-efficacy experiences. By identifying factors associated with the levels of general and task-specific parental selfefficacy, health visitors can gain a fuller appreciation of support needs.

Relevance to practice. To maximise potential through parenting support, attention should be given to addressing factors associated with poorer self-efficacy experiences, including parental tiredness. Equally, practice should be directed at developing community environments that offer exposure to positive praise and the opportunity to practice new skills without facing criticism.

Key words: child discipline, health visiting, parenting, self-efficacy, universal services

Accepted for publication: 18 December 2011

Introduction

Family life and parenting practices are critical factors impacting on the growth and development of young children (Center on the Developing Child at Harvard University 2010). Support for families, particularly in the early years, has been internationally identified as a priority for improving a nation's health and not least, its economic prosperity

Authors: Karen A Whittaker, RN, RHV, BN(Hons), MSc, PGCE, 1,2 PhD, Xxxxx, School of Health, University of Central Lancashire, Preston; Sarah Cowley, RN, RHV, HVT, BA (Hons), PhD, PGDE, Professor, King's College, London, UK

Correspondence: Karen A Whittaker, School of Health, University of Central Lancashire, Preston PR1 2HE, UK.

Telephone: +1772 893786. E-mail: kwhittaker1@uclan.ac.uk (Heckman & Masterov 2007, Mustard 2007). Programmes designed to improve early life experiences often aim to impact on parenting self-efficacy, as a means of encouraging positive parenting practices (Gross *et al.* 2003, Sanders & Woolley 2005). In England, support for parents has been made available through a variety of means, and currently, the health visiting workforce is particularly being invested in to deliver, locally, a continuum of services for families (Department of Health [DH] 2011). In this paper, findings from a survey of self-efficacy experiences completed as part of a larger realistic evaluation of parenting support are drawn on to discuss how practising health visitors' seeking to enhance family well-being needs to act opportunistically to influence potential sources of self-efficacy present in communities. The

Background

The organisation of support for parents

health nurse and public health nurse.

In England, the Department of Health (2011) indicates that families should be able to access different levels of support including help with parenting from health visiting services in convenient local settings as well as the home. Health visitor practice described in a 'new service model' (DH 2011) involves engaging with informal community networks that can provide a foundation of support for all parents and simultaneously leading the universally available Healthy Child Programme (DH 2009). This puts all parents in touch with formal healthcare services relevant to supporting early life (universal level service) and provides an opportunity for identifying when families have either short-term additional needs (universal plus level service) or complex needs that warrant focused help over a longer period (universal partnership plus level service).

UK term health visiting will be used throughout to represent

titles for similar roles elsewhere, such as child and family

For parents, social support often exists as a complex web of formal and informal sources of help (Ghate & Hazel 2002). Importantly, different sources can have a complementary function extending social networks and availability of help (Cochran & Henderson 1990, Edwards & Gillies 2005), but equally, may provide contradictory messages and undermine what might otherwise be helpful sources of influence (Whittaker & Cowley 2010). With this complexity, support for parents is often identified by the mode of contact, hence described as home visiting, group or increasingly, electronic media support.

Formal contact in the home for parents of preschool children has been provided on a universal basis by health visitors in several industrialised nations both in Europe (Kamerman & Kahn 1993, Skovgaard et al. 2005, Cowley et al. 2007) and elsewhere (Wilson 2001, Briggs 2006, Williams et al. 2008). Alternatively, home visitor support may only be offered on a selective basis to high-risk groups, such as teenage parents, as is more common in the United States (Council on Community Pediatrics 2009). International evidence now shows that sustained home visiting enables parents to develop trusting relationships with practitioners (Olds-2007, Appleton & Cowley 2008, Kardama- 3 nidis et al. 2009). Practices based on principles of respect, as for example recommended by the 'Family Partnership Model (FPM)' (Davis & Day 2010), enable parents to disclose sensitive information and visitors to be responsive to parent's needs (Hogg & Worth 2009). Practitioners therefore, through home contact, have the opportunity to develop parents' personal resources, such as self-esteem and selfefficacy.

By contrast, group contact provides support through structured parenting training programmes or more informally through parent-child community groups. In England, both forms of support are universally available via local public bodies known as Children's Centres (DH 2009). The structured programmes delivered by trained facilitators, including health visitors, offer parents preventive guidance by focusing on understanding and responding to children's needs (Barlow & Underdown 2005). By learning appropriate discipline practices, parents help children develop selfcontrol (Stewart-Brown 2008), an important factor for managing later risk (Moffitt et al. 2011). Informal groups, generally led by lay or paraprofessional workers, aim principally to bring parents and children together to socialise and play, thus learning emerges as a by-product of the social experience.

The remaining mode of contact is via media methods. This increasingly means universal provision of information and peer support through discussion boards at dedicated internet sites (Long 2004). In the UK, social networking sites like mumsnet.com offer peer support, with netmums.com, operating an additional professional service, with health visitors hosting an online drop-in clinic (Netmums 2010a).

Social learning as a theoretical basis for parenting interventions

An early review identified that successful parenting programmes were generally underpinned by the theoretical principles of 'modelling', 'behavioural rehearsal' and 'positive reinforcement feedback' (O'Dell 1974). These principles, along with emotional arousal, translate as vicarious

experiences, performance mastery and verbal persuasion, as sources of influence on beliefs about personal capabilities or self-efficacy, a component of social learning theory (Bandura 1997). Contemporary parenting interventions continue to be underpinned by social learning principles (Sutton et al. 2004, Olds et al. 2007), and improvements to parents' personal resources, as indicated by higher parenting efficacy, are regarded as a viable outcome from programmes such as the 'Incredible Years' (Gross et al. 2003, Kim et al. 2008), 'Triple-P Positive Parenting' (Sanders & Woolley 2005) and the nurse-family partnership (Olds et al. 2007). This body of research, however, offers little discussion on the sources of influence that generated improved parenting self-efficacy. Furthermore, self-efficacy theory, although identified as pertinent to health visiting practice (Bloomfield et al. 2005, Whittaker & Cowley 2006, Bloomfield & Kendall 2010), has been little used in this field.

Research suggests that parents' self-efficacy beliefs can be central to their parenting practices (Bloomfield *et al.* 2005, Sanders & Woolley 2005). With poor parental self-efficacy, there is an increased susceptibility towards learned helplessness and thus a poor motivation to address difficulties (Donovan *et al.* 1990). Furthermore, it is a fragile experience that can be disrupted by maternal mood, particularly postnatal depression (Zayas *et al.* 2005, Weaver *et al.* 2008), and can be enhanced by stronger informal social support (Leahy-Warren *et al.* 2014).

Evidence for parenting interventions

A variety of reviews questioning evidence for parenting intervention efficacy are available (Woolfenden et al. 2002, Barlow et al. 2005, Dretzke et al. 2009), but they provide an incomplete picture for those developing universal preventive services. First, despite knowing that group parenting interventions can be effective in having a positive impact on parental well-being, less certainty can be expressed about the factors involved in outcome achievement (Barlow et al. 2003, Hastings & Beck 2004). A second concern is that primary studies examining parenting interventions typically involve parents with very high-level needs associated with caring for children with behaviour disorders. Less attention has been given to the larger population of parents accessing support available at the universal end of the service provision continuum. The purpose of the research reported here was to identify the common experience of parental self-efficacy for those accessing universal parenting services. This work was completed as the first part of a larger realistic evaluation (Pawson & Tilley 1997) concerned with understanding how a parenting programme worked and achieved outcomes.

Methods

Design

A descriptive survey to establish the context for service provision was used.

Study methods

The first part of the study started with the development and pilot test of a questionnaire incorporating outcome measures for parental self-efficacy. The questionnaire survey was targeted at adult users of a multi-component parenting support service available in two geographical wards in a North of England city. For the most part, parents self-referred to service drop-in groups and parenting courses or were recommended by health visitors to receive additional home visiting. The self-referral option created uncertainly about exact numbers accessing services, so non-probability purposive and chunk sampling methods (Aday & Cornelius 2006) were used to include all adults using the parenting service over nine months.

Ethics and governance

Permission to proceed with the realistic evaluation was granted by an NHS Research Ethics Committee and the NHS organisation hosting the parenting service. Guidance with field management of the research came from a stakeholder advisory group consisting of five local parents, two practitioners, three University academics and a regional representative from the charity, Parenting-UK. The group met regularly and debated a range of study issues including ethical practice, questionnaire and information sheet design, field access, interpretation of results and dissemination.

The survey questionnaire

The survey questionnaire (available from the first author) was piloted over a period of six months with assistance from advisory group members who reviewed early drafts and debated results that have already been reported (Whittaker & Cowley 2006). Three differently coloured versions were designed for parents either caring for a baby (<13 months), a toddler (1–4 years) or a school child (5–11 years). Illustrations were included to improve the visual appeal of the questionnaire and divided it into sections containing self-efficacy scales, questions about sources of self-efficacy and demographic details.

Self-efficacy scales

The Bandurian position that assessment of self-efficacy requires a focus on a domain of functioning (Bandura 2006) was adopted, which led to the selection of parenting domain measures rather than measures for general self-efficacy. Those used were the Parenting Self-Agency Measure (PSAM) (Dumka et al. 1996) and subscales from the Self-Efficacy for Parenting Tasks Indexes (SEPTI) toddler and school versions (Coleman & Karraker 2000, 2003). The PSAM, a five-itemed measure, requires the respondent to identify on a 5-point Likert scale how often they felt or thought like each statement representing the domain of parenting. Response categories range from never = 1 - always = 5, with lower scores indicative of lower parental self-efficacy. For the toddler (SEPTI-TS) and school child (SEPTI), only three subscales with content most relevant to the service parenting course were selected for the questionnaire. All the items were rated on a six-point likert scale with possible responses ranging from strongly agree to strongly disagree, with higher scores being indicative of higher self-efficacy. Satisfactory internal consistency had been previously established by the authors of the scales, and the alpha coefficient score results were at acceptable levels when tested with the UK pilot sample. These were as follows: PSAM = 0.76; SEPTI-TS subscales for discipline = 0.66 (improved to 0.79 with questions 4 and 7 removed), play = 0.84 and routine = 0.7; SEPTI School subscales for discipline = 0.86, recreation = 0.88 and nurturance = 0.58 (Whittaker & Cowley 2006).

Self-efficacy source questions

The second section focused on the four main sources of self-efficacy identified by Bandura (1997). This component was included to identify parent exposure to sources of influence that may impact on parenting self-efficacy experiences. Two questions were applied to each of the four known sources of self-efficacy, mastery of experience, vicarious experience, verbal persuasion and physical/psychological state. Each pair of source questions, with the exception of the vicarious experience source, contained a negative and positive statement to reflect the double-sided nature of these sources. Participants were asked to identify how often they had had this experience using an ordinal scale with the categories 'often', 'sometimes', 'rarely' and 'never'. For analytical purposes, these were later coded from 104, with 1 = 'often' and 4 = 'never'.

Data analysis

A log of questionnaire return was kept using a Microsoft Excel spreadsheet, and questions were coded and inputted

statistics were computed to describe the sample. Inferential statistics were used to check for relationships between self-efficacy scores, demographic variables and questions about sources of self-efficacy. Specifically, the internal reliability of PSAM and SEPTI subscales was assessed by computing alpha coefficient scores. The unrelated *t*-test was used to assess self-efficacy mean scores against particular demographic variables, and a one-way analysis of variance (one-way ANOVA) was applied to test for differences between categories of parent participants (parents of children who were babies, toddlers and school aged) all completing the PSAM. To assess for relationships between the self-efficacy source questions and scale scores, correlation coefficients were computed.

Survey administration

Advance notice of the survey was provided through the study advisory group and field researcher visits to each of the health visiting teams in the designated geographical area. To improve study participant recruitment advisory group, parent stakeholders and practising health visitors assisted the field researcher in distributing study information sheets and a total of 278 questionnaires at parenting drop-in groups, parenting courses and the start of a period of health visitor one-to-one parenting support. Parents were asked to complete the questionnaire thinking about one particular child. The survey objective and instructions for completion were detailed on the questionnaire front sheet. They could be completed anonymously, although participants were asked to sign a separate consent form. In total, 168 questionnaires were returned, achieving a sixty per cent response rate. Of these, 62 (37%) had been distributed by professionals, 56 (33%) by parents and 50 (30%) by the researcher.

Results

Demographic details of the study participants provide an insight into who was accessing the parenting service, whilst results from the scales are used to examine parenting self-efficacy experiences and exposure to sources of influence.

Demographic features

Forty-five per cent (n = 75) returned toddler questionnaires, 35% (n = 59) baby and 20% (n = 34) school-aged questionnaires. As shown in Table 1, most of the survey participants, and therefore people accessing the parenting service, were women, were not in paid employment and half were aged between 21–30 years. A sizeable proportion of participants

Table 1 Demographic details for participants

	Frequency, $n = 168$		
Demographic features	n (%)	Missing, n (%	
Gender – female	159 (95)	_	
Age			
16-20 years	19 (11)	3 (2)	
21–30 years	83 (49)	3 (2)	
31–40 years	57 (34)	3 (2)	
41–50 years	3 (2)	3 (2)	
≥51 years	1 (0.5)	3 (2)	
Have a partner	141 (84)	2 (1)	
Lives with partner	124 (74)	1 (0.6)	
Lone adult household	36 (21)	1 (0.6)	
Currently working	40 (24)	10 (6)	
Minority ethnic origin	78 (46)		
Non-minority ethnic origin	91 (54)	_	
Services accessed at point of questionna	ire completion	on	
Parenting training courses	84 (50)	_	
(positive parenting training,			
baby massage training)			
Drop-in groups (parent and toddler,	51 (31)	_	
teenagers parents group, dads group)			
Health visiting services (baby clinics,	19 (11)	_	
home visits)			
Not stated	14 (8)	_	
An additional service accessed	105 (62.5)	17 (10)	

(n = 105, 62.5%) were making use of more than one form of parenting support and those accessed at the point of questionnaire completion included formalised courses (n = 84, 50%) and individual contact (n = 19, 11%) as well as drop-in groups (n = 51, 31%).

Almost half of the participants (n = 78, 46%) were of minority ethnic origin and many of whom were of South Asian descent. The average household size was four people,

but this varied between 2–11 people. When cross-tabulated with ethnicity and age, it was apparent that those living in lone adult households (n = 36) were typically white British participants (n = 29) ($\chi^2 = 13.42$, df = 1, p < 0.001) and tended to be aged 16–25 years (n = 20) ($\chi^2 = 8.787$, df = 1, p = 0.012). The mean number of children (aged 0–16 years) living with participants was two.

Parenting experiences of self-efficacy

Reliability testing with the whole sample produced a Cronbach's alpha coefficient of 0.67 for the PSAM. Similarly, sufficiently internal reliability was also found for the toddler and school versions of the SEPTI scales with Cronbach's alpha coefficients calculated for toddler subscales: discipline (five items) = 0.87; play (seven items) = 0.85; routine (six items) = 0.76 and for school subscales: discipline (eight items) = 0.84; recreation (seven items) = 0.75; nurturance (seven items) = 0.61.

The PSAM was the only scale to be completed by all participants (Table 2), and the one-way anova result indicated that there was no significant difference in PSAM scores between the baby, toddler and school groups [F(2, 166) = 0.628, p = 0.535]. The mean and median PSAM results suggest a good level of parenting self-efficacy across all participants although it appeared that there were some small demographic differences. Specifically, the unrelated t-test indicates differences in response to minority (mean = 21) and non-minority (mean = 20) ethnic origin parents (t = -2.23, df = 162, p = 0.027). Differences were also found depending on whether the respondent lived with their partner (mean = 21) or not (mean = 20) (t = 2.22, df = 162, p = 0.028).

Table 2 Mean PSAM, SEPTI-TS and SEPTI-school scores

Measure	N	Minimum	Maximum	Median 95% Confidence interval		Mean	SE
PSAM							
All parents	166	1	25	21	20-20-9	20	0.20
Parents with							
Child < 13 months (baby)	59	15	25	20	19-6-20-9	20	0.35
Child 1-4 years (toddler)	73	13	25	21	19-9-21	20	0.29
Child 5-11 years (school)	34	15	25	21	20-21.8	21	0.45
SEPTI-TS							
Discipline	71	6	30	18	16.9–19.9	18	0.76
Play	72	12	42	35	32·4–35·5	34	0.77
Routine	72	12	36	28	26-28.8	27	0.70
SEPTI-School							
Discipline	32	20	48	34	31.9-37.6	35	1.4
Recreation	32	12	41	34	29·1–33·8	31	1.16
Nurturance	33	28	42	39	36·6–39·2	38	0.63

PSAM, Parenting Self-Agency Measure; SEPTI, Self-Efficacy for Parenting Tasks Index.

The advantage of including the domain-specific SEPTI subscales was that experiences related to specific aspects of parenting could be considered. The results (Table 2), with the exception of those for the toddler discipline subscale, indicated good levels of parenting self-efficacy across specific areas of parenting toddlers and school-aged children. No relationships were found between the demographic variables and any of the SEPTI-school subscales, but some weak and inversely related relationships were found with the toddler subscales. These were for the play subscale scores (n = 67) with household size (r = -0.25, p = 0.04) and with number of children (r = -0.26, p = 0.03) and for routine subscale scores (n = 68) with the number of adults in the household (n = -0.35, n = 0.003).

Parenting experience of the sources of self-efficacy

To examine whether there was any relationship between participants sense of self-efficacy and exposure to different community experiences, the PSAM scores were correlated with the questions aimed at exploring the sources of selfefficacy. To accommodate the ordinal level of measurement used here, the nonparametric Spearman rho statistic was calculated (Hicks 1990). The results detailed in Table 3 show very weak relationships, principally with sources concerned with feeling successful, criticised and tired (questions 1, 6 and 7). In addition, different efficacy experiences were suggested by relationships between the source questions. Here, 'feeling tired' positively correlated with often 'giving-in' and negatively correlated with 'often feeling successful in efforts', although again, these relationships were very weak.

Only one of the source questions (feeling tired) showed a weak relationship with the school discipline subscale (n = 32) ($r_s = 0.388$, p = 0.028). Stronger relationships were found between SEPTI-TS subscale scores and some self-efficacy source questions (Table 4). It appeared that 'feeling successful in parenting efforts' did to some degree correlate with higher SEPTI-TS scores for routine, discipline and to a lesser degree play. Opposite to this, those who often felt they 'gave-in' to their child's demands had lower routine and discipline scores. 'Feeling tired' accounted for some scores across all of the subscales, and this source seemed more influential than 'having patience'.

Table 3 Correlation between PSAM scores and self-efficacy sources questions

	PSAM score,	Source questions, $n = 164$						
Spearman's rho (r_s)	n = 164	1	2	3	4	5	6	7
1 Feel successful in efforts as a parent	-0.259**							
2 Feel you 'give in' to child's demands	0.152	0.166*						
3 See someone successfully calm child	-0.035	-0.010	-0.017					
4 See someone successfully teach child	-0.151	0.082	0.030	0.385**				
5 Receive positive comments	-0.029	-0.006	-0.090	0.270**	0.163*			
6 Receive negative comments	0.235**	-0.208**	0.144	0.196*	0.043	0.139		
7 Feel too tired to deal with difficult behaviour	0.257**	-0.285**	0.308**	-0.021	-0.007	-0.165*	0.207**	
8 Have the patience to deal with difficult behaviour	· -0·186*	0.134	0.073	0.024	-0.001	0.118	-0.104	-0.204**

PSAM, Parenting Self-Agency Measure.

Table 4 SEPTI-TS correlations with self-efficacy sources

	SEPTI-TS subscales, Spearman's rho (r _s)					
Self-efficacy source questions	Discipline, $n = 70$	Play, <i>n</i> = 71	Routine, $n = 71$			
Feel successful in efforts	-0.408***	-0.298*	-0.559***			
Often 'giving-in'	0.315**	0.147	0.513***			
Given negative comments	0.197	0.224	0.243*			
Feel too tired	0.429**	0.38**	0.434***			
Feel do have patience	-0.341**	-0.379**	-0.215			

SEPTI, Self-Efficacy for Parenting Tasks Index.

^{*}Correlation is significant at the 0.05 level (two-tailed).

^{**}Correlation is significant at the 0.01 level (two-tailed).

^{*}p < 0.05, ** $p \le 0.01$, *** $p \le 0.001$.

Discussion

In the background to this paper, the nature of parenting support and health visitor involvement was set out. The rationale for this area of public health activity rests on the need for early investment in the lives of children, even in developed societies. Indeed, action to support families, particularly parents, to meet the challenges of raising children is identified as essential for working towards strengthening population health and economic prosperity (Mustard 2007).

To assist families, there are various modes of parenting support, which commonly aim to strengthen parental self-efficacy and competence. This study identified women as primary users of universal parenting support and was interested in how self-efficacious they felt as parents and the contextual factors that could be shaping these experiences. The findings are consistent with arguments that several contextual factors are important to personal self-efficacy experiences and provide several indications of the relevance of self-efficacy theory to contemporary health visiting practice.

Common parental self-efficacy experiences

In exploring common experiences of parental self-efficacy, it was apparent that parenting needs might not always be what they seem and hence personalised assessment is as important in health visiting, as it is in others spheres of nursing. This is illustrated in data concerning living situations, where participants from larger households tended to have poorer self-efficacy for managing toddler routines or offering play opportunities. That is, in households with more adults and theoretically carers, the parental load was not necessarily lightened. This challenges assumptions that those in lone parent households are the most vulnerable and marks out the need for practitioners to seek a complete picture when defining needs for additional support.

Examination of self-efficacy scores in the specific domains of parenting also shows how initial impressions can be misleading. Here, the difference in the domain general PSAM and task-specific SEPTI scale totals suggests that whilst parents experienced relatively good general parental self-efficacy, they were less sure of their capability in particular areas. Specifically, less confidence was expressed in managing child discipline. Furthermore, examination of experiences acting as sources of influence on self-efficacy showed that those who often 'felt tired' or who 'gave-in' to their child's demands indicated lower self-efficacy in managing discipline and routine. Similarly, parents expressed greater self-efficacy in managing discipline and routine, if they identified them-

selves as often 'feeling successful in their efforts as a parent'. Thus, parents had poorer self-belief in managing specific child rearing challenges when they were physically compromised by tiredness, or had, because of giving-up, rarely experienced positive mastery, whereas those experiencing mastery and success were more positive about their capabilities in the areas of discipline and routine. These results not only highlight important areas for attention by health visitors, but are consistent with Bandura's (1997) messages about the particularly powerful influence of being able to master a task and equally the damaging effects of the reverse; repeated exposure to non-mastery and giving-up.

Managing discipline appropriately, including establishing routines, is an important component of early child rearing patterns because through this practice parents can help their children develop self-control; a fundamental skill for later life (Moffitt *et al.* 2011). The challenge for parents is to be consistent with their positive discipline practices and to avoid 'giving-in' when they feel more stressed, as frequent inconsistencies in parenting practices (a lax style) have been associated with greater childhood behaviour difficulties (Ang 2008).

Factors Bandura (1997) identified as impacting on the ability to persist in mastering a new skill or means of managing a difficulty, such as child behaviour, include the individual's physical and psychological status, as well as feedback received from others. Here, parents' experiences of 'feeling tired' correlated positively with the belief that they often 'received negative comments' and often 'gave-in' to their child's demands. This reduced exposure to helpful sources perhaps explains why some of those who expressed 'feeling tired' were more inclined to have lower parenting self-efficacy scores.

Opportunities for strengthening parenting support

To strengthen support services for parents, practitioners, such as health visitors, face a fourfold challenge. This includes knowing individual families sufficiently to appropriately understand their requirements regarding the levels of self-efficacy experienced and concurrent exposure to sources of influence. Next, they need to be ready to provide guidance relevant to key aspects of parenting where parents express least self-belief; do this in a sensitive and balanced manner to avoid undermining any sense of general parenting self-efficacy; and use methods that can reverse exposure to negative and build on positive sources of influence. Achieving this, however, is likely to depend on the ability to have enough personal contact with each family. Contact as part of the universal component of a comprehensive child health

232425

programme, such as the Healthy Child Programme in England (DH 2009), is important for two reasons. It offers a basis for determining need and identifying who might benefit from the more intensive modes of support and thereby provides data required for planning resource allocation from the care continuum.

The finding with respect to tiredness is one of the notes given that this can feature as part of the transition to parenthood (Barclay et al. 1997). In a survey of over 5000 mothers using online parenting support, 50% of respondents cited lack of sleep as one of their main day-to-day stressors (Netmums 2010b). Persistent tiredness can contribute to low mood and possible depressive illness (Sharpe & Wilks 2002), which can disrupt parental self-efficacy (Zayas et al. 2005, Weaver et al. 2008). Moreover, arguments for taking parental tiredness seriously are supported by evidence that poorer self-efficacy adversely affects maternal sensitivity (Donovan et al. 2005) whilst attentive sensitive parenting enables positive infant neurological development (Gerhardt 2004). Rather than minimising tiredness as an automatic parenting experience, parents' felt experiences can be acknowledged. This would create the opportunity to address the experience directly and introduce solution-focused approaches for managing parents' sleep/wakefulness difficulties. The intention was, to minimise threats that tiredness could be posing to parental self-efficacy and sensitive parenting practices.

If practitioners understand the areas where parents feel least self-efficacy, they can maximise exposure to positive sources of influence available through universal childhood services. This includes making the most of contact in the home, a support method valued by parents if they are able to establish firm relationships with health visitors (Hogg & Worth 2009). Kardamanidis et al.'s (2009) Australian study showed how the FPM (Davis & Day 2010), formerly known as 'parent advisor approach' (Davis et al. 2002), enables respectful relationships to develop, which can be used to facilitate a shared exploration of parental need and identity of tasks with which parents feel least self-efficacious. Part of the exploration would include identifying the different sources of support and influence in families and neighbourhoods (real or virtual), determining whether these have the potential to contribute positively or negatively to parenting self-efficacy experiences. By listening to parents' concerns, then guiding them in the process of identifying their own solutions, practitioners would enable investment in parents' own personal resources and thus psychological sources of influence on self-efficacy.

A broad-based preventive service, such as that described in the 'new service vision' for English health visitors (DH 2011), provides several opportunities for action directed at maximising exposure to positive sources of self-efficacy or active management of potential negative exposures. The universal service component might include home contact as described above, along with practitioner involvement in community parent-child groups. In community situations, practitioners have a role in facilitating respectful communication between participants, encouraging exposure to positive verbal persuasion messages and role modelling in community situations. Where parents have additional self-efficacy needs, parenting training programmes that incorporate role play and feedback would provide scope for task-specific role modelling and the chance to practice new parenting strategies in a safe situation. From this, an opportunity arises for exposure to positive verbal persuasion messages along with vicarious and mastery experiences. What is more, the increasing use of social networking internet sites and specifically the growth in those aimed at parents, would suggest that virtual communities are also environments where practitioners could have an important role to play in shaping exposure to self-efficacy sources. In this regard, the example of Netmums employing health visitors to support online clinics and parenting programmes might prove to be a useful model for practice capable of complementing other modes of universal service provision.

Limitations

The survey provided a one-off measure, a snapshot at a given period of time, of common parental self-efficacy experiences for users of any one of several parenting support facilities. This meant it would not have been appropriate, in this instance, to repeat the questionnaire to assess for change in outcome measure results. This was because the contextual situations for each questionnaire completion in a multi-component service were likely to vary between measurements. What is more, parents used several facilities simultaneously or at different times, so it was difficult to define a clear start and end point to use of formal parenting support. This meant that the survey data did not necessarily represent the true starting point of service use, so reliable measurement of changes to common parenting self-efficacy experiences was not feasible in this instance.

However, the outcome measures used in the study could be used elsewhere as valid and reliable indicators of parenting programme success, that is, in a study situation where greater assurances could be made about the intervention being measured and where a level of consistency in circumstances could be achieved for each questionnaire completion. The outcome measure Cronbach's alpha coefficients would, however, suggest that the SEPTI-TS and the SEPTI school subscales, with the exception of the nurturance subscale,

were more stable as measures of parental self-efficacy than the PSAM. These results and the fact that the PSAM was a general domain measure as opposed to a task-specific measure also provide some explanation for why only weak correlations were found between SEPTI-TS and the PSAM scale results.

Conclusion

The paper started by highlighting how support for parents has become a recognised feature of public services aimed at improving the life chances of children. The ways support can be organised have been outlined, drawing attention to the move in England to make services available as part of a broad-based preventive model whereby those with greater need have access to greater support (DH 2011). This background was presented as a basis for the discussion of findings from the first part of a realistic evaluation study; specifically survey evidence of parental self-efficacy experiences for those accessing universal support services. Survey findings indicated that parents generally experienced good levels of parental self-efficacy, although evidence from the task-specific scales would suggest that managing discipline was an area of parenting with which they felt least efficacious. There was also some suggestion of relationships existing between poorer self-efficacy and factors such as feeling tired, receiving negative comments and often 'givingin' to a child's demands. These findings have implications for providers of universal child health programmes and in particular health visitors who are able to make contact with families in a wide range of community situations. Knowledge of social learning theory and understanding its application by managing exposure to sources of influence for self-efficacy development can provide practitioners with a valuable means of helping. Moreover, by developing an understanding of parental self-efficacy experiences in a community, practitioners have the opportunity to strengthen local parenting support through practices that are consistent with the development of environments rich in positive sources of

self-efficacy. Practice of this kind could have implications for parent-child relationships and thereby the longer-term development of children and the communities in which they live

Relevance to clinical practice

- Four sources of influence are known to be important to the strength of self-efficacy, which are mastery of experience, vicarious experience, verbal persuasion and physical/psychological state.
- Parents experience of 'feeling tired', 'receiving negative comments' and 'often giving-in to child's demands' are particularly associated with lower perceived parental selfefficacy – these are experiences that health visitors and colleagues should take seriously when delivering universal child health services
- Practitioners should be alert to the possibility that even when parents demonstrate good general levels of parental self-efficacy, they may be less sure of their capabilities in certain areas of parenting, such as discipline.

Acknowledgements

We are grateful to all parents who gave time to complete the questionnaire survey, those contributing to the study advisory group, and Dr Chris Sutton who provided statistical guidance. The Department of Health is thanked for funding the NCC RCD doctoral research training fellowship (grant number RDO/33/89).

Contributions

Study design: KW, SC; data collection and analysis: KW and manuscript preparation: KW, SC.

Conflicts of interest

The authors have no conflict of interest.

References

Aday LA & Cornelius LJ (2006). A Comprehensive Guide. Designing and Conducting Health Surveys, 3rd edn. Jossey-Bass, San Francisco, CA.

Ang RP (2008) Dysfunctional parenting behaviors and parenting stress among mothers of aggressive boys. *Child & Family Behavior Therapy* 30, 319–336.

Appleton JV & Cowley S (2008) Health visiting assessment – unpacking critical attributes in health visitor needs assessment practice: a case study. International Journal of Nursing Studies 45, 232–245.

Bandura A (1997) Self-Efficacy: The Exercise of Control. W.H. Freeman and Company, New York, NY.

Bandura A (2006) Guide for constructing self-efficacy scales. In *Self-Efficacy Beliefs of Adolescents* (Pajares F & Urdan T eds). Information Age Publishing, Greenwich, CT, pp. 307– 337.

Barclay L, Everitt L, Rogan F, Schmied V & Wyllie A (1997) Becoming a mother – an analysis of women's experience of

- early motherhood. Journal of Advanced Nursing 25, 719-728.
- Barlow J & Underdown A (2005) Promoting the social and emotional health of children: where to now? The Journal of the Royal Society for the Promotion of Health 125, 64-70.
- Barlow J, Coren E & Stewart-Brown SSB (2003) Parent-training programmes for improving maternal psychosocial health. Cochrane Database of System-Reviews 2003. Art. CD002020.
- Barlow J, Parsons J & Stewart-Brown S (2005) Preventing emotional and behavioural problems: the effectiveness of parenting programmes with children less than 3 years of age. Child: Care Health Development 31, 33-42.
- Bloomfield L & Kendall S (2010) Audit as evidence: the effectiveness of '123 Magic' programmes. Community Practitioner 83, 26-29.
- Bloomfield L, Kendall S, Applin L, Attarzadeh V, Dearnley K, Edwards L, Hinshelwood L, Lloyd P & Newcombe T (2005) A qualitative study exploring the experiences and views of mothers, health visitors and family support centre workers on the challenges and difficulties of parenting. Health & Social Care in the Community 13, 46-55.
- Briggs C (2006) Nursing practice in community child health: developing the nurse-client relationship. Contemporary Nurse: A Journal for the Australian Nursing Profession 23, 303-311.
- Center on the Developing Child at Harvard University (2010) The Foundations of Lifelong Health Are Built in Early Childhood. Available at: http://www. developingchild.harvard.edu (accessed 23 September 2011).
- Cochran M & Henderson CR (1990) Formal supports and informal social ties: a case study. In Extending Families: The Social Networks of Parents and Their Children (Cochran M, Larner M, Riley D, Gunnarsson L & Henderson CR eds). Cambridge University Press, Cambridge, pp. 230-261.
- Coleman PK & Karraker KH (2000) Parenting self-efficacy among mothers of school-age children: conceptualization, measurement and correlates. Family Relations 49, 13-24.
- Coleman PK & Karraker KH (2003) Maternal self-efficacy beliefs, compe-

- tence in parenting and toddlers' behavior and developmental status. Infant Mental Health Journal 24, 126-
- Council on Community Pediatrics (2009) The role of preschool home-visiting programs in improving children's developmental and health outcomes. Pediatrics 123, 598-603.
- Cowley S, Caan W, Dowling S & Weir H (2007) What do health visitors do? A national survey of activities and service organisation. Public Health 121, 869-879.
- Davis H & Day C (2010) Working in Partnership with Parents, 2nd edn. Pearson, London,
- Davis H, Day C & Bidmead C (2002) Working in Partnership with Parents. The Psychological Corporation, Lon-
- Department of Health (2009) Healthy Child Programme: Pregnancy and the First Five Years of Life. DH Publications,
- Department of Health (2011) Health Visitor Implementation Plan 2011-15. Department of Health, London.
- Donovan WL, Leavitt LA & Walsh RO (1990) Maternal self-efficacy: illusory control and its effect on susceptibility to learned helplessness. Child Development 61, 1638-1647.
- Donovan WL, Leavitt LA & Taylor N (2005) Maternal self-efficacy and experimentally manipulated infant difficulty effects on maternal sensory sensitivity: a signal detection analysis. Developmental Psychology 41, 784-798.
- Dretzke J, Davenport C, Frew E, Barlow J, Stewart-Brown S, Bayliss S, Taylor R, Sandercock J & Hyde C (2009) The clinical effectiveness of different parenting programmes for children with conduct problems: a systematic review of randomised controlled trials. Child and Adolescent Psychiatry and Mental Health 3, 7.
- Dumka LE, Stoerzinger HD, Jackson KM & Roosa MW (1996) Examination of the cross-cultural and cross-language equivalence of the parenting self-agency measure. Family Relations 45, 216-222.
- Edwards R & Gillies V (2005) Resources in Parenting: Access to Capitals Project Report. London South Bank University, London.

- Gerhardt S (2004) Why Love Matters: How Affection Shapes a Baby's Brain. Routledge, Hove, East Sussex.
- Ghate D & Hazel N (2002). Parenting in Poor Environments. Stress, Support and Coping. Jessica Kingsley Publishers, London.
- Gross D, Fogg L, Webster-Stratton C, Garvey C, Julion W & Grady J (2003) Parent training of toddlers in day care in low-income urban communities. Journal of Consulting & Clinical Psvchology 71, 261-278.
- Hastings RP & Beck A (2004) Practitioner review: stress intervention for parents of children with intellectual disabilities. Journal of Child Psychology & Psychiatry 45, 1338-1349.
- Heckman JJ & Masterov DV (2007) The productivity argument for investing in young children. Applied Economic Perspectives and Policy 29, 446-493.
- Hicks CM (1990). Research and Statistics. A Practical Introduction for Nurses. Prentice Hall, London.
- Hogg R & Worth A (2009) What support do parents of young children need? A user-focused study. Community Practitioner 82, 31-33.
- Kamerman SB & Kahn AJ (1993) Home health visiting in Europe. 3. Available at: http://futureofchildren.org/futureofchildren/publications/journals/archive.xml (accessed 30 April 2011).
- Kardamanidis K, Kemp L & Schmied V (2009) Uncovering psychosocial needs: perspectives of Australian child and family health nurses in a sustained home visiting trial. Contemporary Nurse: A Journal for the Australian Nursing Profession 33, 50-58.
- Kim E, Cain KC & Webster-Stratton C (2008) The preliminary effect of a parenting program for Korean American mothers: a randomized controlled experimental study. International Journal of Nursing Studies 45, 1261-
- Leahy-Warren P, McCarthy G & Corcoran P (2014) First-time mothers: social support, maternal parental self-efficacy and postnatal depression. Journal of Clinical Nursing ???, ???-???, doi: 10.1111/j.1365-2702.2011.03701.x.
- Long N (2004) e-Parenting. In Handbook of Parenting. Theory and Research for Practic (Hoghughi M & Long N eds). London, Sage, pp. 369-379.

- Moffitt TE, Arseneault L, Belsky D, Dickson N, Hancox RJ, Harrington H, Houts R, Poulton R, Roberts BW, Ross S, Sears MR, Thomson WM & Caspi A (2011) A gradient of childhood self-control predicts health, wealth and public safety. *Proceedings of the National Academy of Sciences* 108, 2693–2698.
- Mustard JF (2007) Experience-based brain development: scientific underpinnings of the importance of early child development in a global world. In *Early Child Development from Measurement to Action. A Priority for Growth and Equity* (Young ME & Richardson L eds). The World Bank, Washington, D.C., pp. 35–63.
- Netmums (2010a) A Mum's Life. Available at: http://www.netmums.com/campaigns/A_Mum_s_Life.656/ (accessed 30 April 2011).
- Netmums (2010b) Netmums Parent Supporters. Available at: http://www. netmums.com/h/n/SUPPORT/drop_in_ clinic/ (accessed 30 April 2011).
- O'Dell S (1974) Training parents in behavior modification. *Psychological Bulletin* **81**, 418–433.
- Olds DL, Sadler L & Kitzman H (2007)
 Programs for parents of infants and toddlers: recent evidence from randomized trials. *Journal of Child Psychology and Psychiatry* 48, 355–391.

- Pawson R & Tilley N (1997) Realistic Evaluation. Sage, London.
- Sanders MR & Woolley ML (2005) The relationship between maternal self-efficacy and parenting practices: implications for parent training. *Child: Care, Health & Development* 31, 65–73.
- Sharpe M & Wilks D (2002) ABC of psychological medicine: fatigue. *British Medical Journal* 325, 480–483.
- Skovgaard AM, Olsen EM, Houmann T, Christiansen E, Samberg V, Lichtenberg A & Jorgensen T (2005) The Copenhagen County child cohort: design of a longitudinal study of child mental health. Scandinavian Journal of Public Health 33, 197–202.
- Stewart-Brown S (2008) Improving parenting: the why and the how. *Archives of Disease in Childhood* 93, 102–104.
- Sutton C, Utting D & Farrington D (2004).

 Support from the Start. Working with
 Young Children and Their Families to
 Reduce the Risks of Crime and Antisocial Behaviour. DfES Publications,
 Nottingham.
- Weaver CM, Shaw DS, Dishion TJ & Wilson MN (2008) Parenting self-efficacy and problem behavior in children at high risk for early conduct problems: the mediating role of maternal depression. *Infant Behavior and Development* 31, 594–605.

- Whittaker KA & Cowley S (2006) Evaluating health visitor parenting support: validating outcome measures for parental self-efficacy. *Journal of Child Health Care* 10, 296–308.
- Whittaker KA & Cowley S (2010) An effective programme is not enough: a review of factors associated with poor attendance and engagement with parenting support programmes. *Children & Society* ???, ???-???, doi: 10.1111/j.1099-0860.2010.00333.x₇
- Williams R, Clinton J & Biscaro A (2008) Ontario and the enhanced 18-month well-baby visit: trying new approaches. *Pediatric Child Health* 13, 850– 856.
- Wilson HV (2001) Power and partnership: a critical analysis of the surveillance discourses of child health nurses. *Journal of Advanced Nursing*. **36**, 294–301.
- Woolfenden SR, Williams K & Peat JK (2002) Family and parenting interventions for conduct disorder and delinquency: a meta-analysis of randomised controlled trials.

 Archives of Disease in Childhood 86, 251–256.
- Zayas LH, Jankowski KRB & McKee MD (2005) Parenting competency across pregnancy and postpartum among urban minority women. *Journal of Adult Development* 12, 53–62.

The Journal of Clinical Nursing (JCN) is an international, peer reviewed journal that aims to promote a high standard of clinically related scholarship which supports the practice and discipline of nursing.

For further information and full author guidelines, please visit JCN on the Wiley Online Library website: http://wileyonlinelibrary.com/journal/jocn

Reasons to submit your paper to JCN:

High-impact forum: one of the world's most cited nursing journals and with an impact factor of 1·228 – ranked 23 of 85 within Thomson Reuters Journal Citation Report (Social Science – Nursing) in 2009.

One of the most read nursing journals in the world: over 1 million articles downloaded online per year and accessible in over 7000 libraries worldwide (including over 4000 in developing countries with free or low cost access).

Fast and easy online submission: online submission at http://mc.manuscriptcentral.com/jcnur.

Early View: rapid online publication (with doi for referencing) for accepted articles in final form, and fully citable.

Positive publishing experience: rapid double-blind peer review with constructive feedback.

Online Open: the option to make your article freely and openly accessible to non-subscribers upon publication in Wiley Online Library, as well as the option to deposit the article in your preferred archive.

Author Query Form

Journal: JOCN

Article: 4074

Dear Author,

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers on the query sheet if there is insufficient space on the page proofs. Please write clearly and follow the conventions shown on the attached corrections sheet. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.

Many thanks for your assistance.

Query reference	Query	Remarks
1	AUTHOR: Please restricts the qualification to three.	D
2	AUTHOR: Please provide designation for Karen A Whittaker.	D
3	AUTHOR: Olds 2007 has not been included in the Reference List, please supply full publication details.	
4	AUTHOR: Please give manufacturer information for SPSS version 10: company name, town, state (if USA), and country.	
5	AUTHOR: Please provide the volume number and page range for reference Leahy-Warren <i>et al.</i> (2011).	
6	AUTHOR: Please provide the volume number and page range for reference Whittaker & Cowley (2010).	



USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION

Required software to e-Annotate PDFs: <u>Adobe Acrobat Professional</u> or <u>Adobe Reader</u> (version 8.0 or above). (Note that this document uses screenshots from <u>Adobe Reader X</u>)

The latest version of Acrobat Reader can be downloaded for free at: http://get.adobe.com/reader/


Once you have Acrobat Reader open on your computer, click on the Comment tab at the right of the toolbar:





This will open up a panel down the right side of the document. The majority of tools you will use for annotating your proof will be in the Annotations section, pictured opposite. We've picked out some of these tools below:



1. Replace (Ins) Tool – for replacing text. Strikes a line through text and opens up a text box where replacement text can be entered. How to use it Highlight a word or sentence. Click on the Replace (Ins) icon in the Annotations Type the replacement text into the blue box that appears. idard framework for the analysis of m icy. Nevertheless, it also led to exoge ole of strateg n f 🥯 * dthreshe nber of comp 08/06/2011 15:58:17 O is that the st which led ofnain compo be level, are exc nc important works on entry by entire M henceforth) we open the 'black h









USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION

5. Attach File Tool – for inserting large amounts of text or replacement figures.



Inserts an icon linking to the attached file in the appropriate pace in the text.

How to use it

- Click on the Attach File icon in the Annotations section
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.

0.10 0.10


6. Add stamp Tool – for approving a proof if no corrections are required.



Inserts a selected stamp onto an appropriate place in the proof.

How to use it

- Click on the Add stamp icon in the Annotations section.
- Select the stamp you want to use. (The Approved stamp is usually available directly in the menu that appears).
- Click on the proof where you'd like the stamp to appear. (Where a proof is to be approved as it is, this would normally be on the first page).






7. Drawing Markups Tools – for drawing shapes, lines and freeform annotations on proofs and commenting on these marks.

Allows shapes, lines and freeform annotations to be drawn on proofs and for comment to be made on these marks..

How to use it

- Click on one of the shapes in the Drawing Markups section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, move the cursor over the shape until an arrowhead appears.
- Double click on the shape and type any text in the red box that appears.



For further information on how to annotate proofs, click on the Help menu to reveal a list of further options:

